


Electrostatics-Potential

- If 1.0 joule of work is required to move 1.0 coulomb of charge between two points in an electric field, the potential difference between the two points is
 - $1.0 \times 10^0 \text{ V}$
 - $9.0 \times 10^9 \text{ V}$
 - $6.3 \times 10^{18} \text{ V}$
 - $1.6 \times 10^{-19} \text{ V}$

- The diagram below represents a positively charged particle about to enter the electric field between two oppositely charged parallel plates.

+++++



The electric field will deflect the particle

- into the page
 - out of the page
 - toward the top of the page
 - toward the bottom of the page
- What is the total amount of work required to move a proton through a potential difference of 100 volts?
 - $1.60 \times 10^{-21} \text{ J}$
 - $1.60 \times 10^{-17} \text{ J}$
 - $1.00 \times 10^2 \text{ J}$
 - $6.25 \times 10^{20} \text{ J}$
 - The diagram below represents two electrons, e_1 and e_2 , located between two oppositely charged parallel plates.

+++++



Compare the magnitude of the force exerted by the electric field on e_1 to the magnitude of the force exerted by the electric field on e_2 .

- Which electrical unit is equivalent to one joule?
 - volt per meter
 - ampere-volt
 - volt per coulomb
 - coulomb-volt
- If 60 joules of work is required to move 5.0 coulombs of charge between two points in an electric field, what is the potential difference between these points?
 - 5.0 V
 - 12 V
 - 60 V
 - 300 V

- In the diagram below, proton p, neutron n, and electron e are located as shown between two oppositely charged plates.

+++++

(p)

(n)

(e)

The magnitude of acceleration will be greatest for the


- neutron, because it has the greatest mass
 - neutron, because it is neutral
 - electron, because it has the smallest mass
 - proton, because it is farthest from the negative plate
- An electron is accelerated through a potential difference of 2.5×10^4 volts in the cathode ray tube of a computer monitor. Calculate the work, in joules, done on the electron. [Show all work, including the equation and substitution with units.]

Electrostatics-Potential

9. Moving 2.5×10^{-6} coulomb of charge from point A to point B in an electric field requires 6.3×10^{-4} joule of work. The potential difference between points A and B is approximately

1. 1.6×10^{-9} V
2. 4.0×10^{-3} V
3. 2.5×10^2 V
4. 1.0×10^{14} V

10. The diagram below represents a source of potential difference connected to two large, parallel metal plates separated by a distance of 4.0×10^{-3} meter.



Which statement best describes the electric field strength between the plates?

1. It is zero at point B
 2. It is a maximum at point B
 3. It is a maximum at point C
 4. It is the same at points A, B, and C.
11. In an electric field, 0.90 joule of work is required to bring 0.45 coulomb of charge from point A to point B. What is the electric potential difference between points A and B?
1. 5.0 V
 2. 2.0 V
 3. 0.50 V
 4. 0.41 V
12. A potential difference of 10 volts exists between two points, A and B, within an electric field. What is the magnitude of charge that requires 2.0×10^{-2} joule of work to move it from A to B?
1. 5.0×10^2 C
 2. 2.0×10^{-1} C
 3. 5.0×10^{-2} C
 4. 2.0×10^{-3} C

13. If 4.8×10^{-17} joule of work is required to move an electron between two points in an electric field, what is the electric potential difference between these points?

1. 1.6×10^{-19} V
2. 4.8×10^{-17} V
3. 3.0×10^2 V
4. 4.8×10^2 V

Base your answers to questions 14 and 15 on the information below.

A proton starts from rest and gains 8.35×10^{-14} joule of kinetic energy as it accelerates between points A and B in an electric field.

14. What is the final speed of the proton?
1. 7.07×10^6 m/s
 2. 1.00×10^7 m/s
 3. 4.28×10^8 m/s
 4. 5.00×10^{13} m/s
15. Calculate the potential difference between points A and B in the electric field. [Show all work, including the equation and substitution with units.]
16. Which is a vector quantity?
1. electric charge
 2. electric field strength
 3. electric potential difference
 4. electric resistance
17. Which object will have the greatest change in electrical energy?
1. an electron moved through a potential of 2.0 V
 2. a metal sphere with a charge of 1.0×10^{-9} C moved through a potential difference of 2.0 V
 3. an electron moved through a potential of 4.0 V
 4. a metal sphere with a charge of 1.0×10^{-9} C moved through a potential difference of 4.0 V